

# Brain-Wide Mapping of Stimulus Induced Variability Quenching Reveals Modularity of Cortical Network

**Submission ID** 3000187  
**Submission Type** Poster  
**Topic** Neuroscience  
**Status** Submitted  
**Submitter** Teppei Matsui  
**Affiliation** University of Tokyo

## SUBMISSION DETAILS

**Presentation Type** Poster Presentation

**Presentation Abstract Summary** Recent studies have revealed that neurons not directly activated by a given sensory stimulus can nevertheless “respond” to the stimulus by showing reduction of trial-to-trial variability in neural activity (“variability quenching”). Determining the extent of cortical neural circuit that responds to a sensory stimulus is important for understanding how cortical network as a whole process sensory information. Here we used mesoscopic brain activity mapping to find that variability quenching occurred in widespread, bilaterally symmetric, cortical network-module. Each network-module showed variability quenching to particular sensory modality. The network-modules identified by variability quenching closely resembled cortical networks identified by spontaneous activity correlation (a.k.a. resting-state functional connectivity). Thus, although only a subset of neurons is activated, sensory stimulation recruits an entire cortical network-module that share correlated spontaneous activity.

**Paper Upload (PDF)** [ccn\\_abstract\\_Matsui\\_etal.pdf](#)

## Co-author Information

\* Presenting Author

First Name	Last Name	Affiliation	E-mail
Teppei *	Matsui *	University of Tokyo	tematsui@m.u-tokyo.ac.jp
Tomonari	Murakami	University of Tokyo	t-mura@m.u-tokyo.ac.jp
Kenichi	Ohki	University of Tokyo	kohki@m.u-tokyo.ac.jp

## Keywords

**Keywords**

mouse brain connectivity and dynamics
---------------------------------------

mesoscopic imaging
--------------------

neural network modeling
-------------------------