

The Evolution of Tactile Letter Representations in Blind Braille Readers

Submission ID 3000316

Submission Type Poster

Topic Neuroscience

Status Submitted

Submitter Santani Teng

Affiliation Massachusetts Institute of Technology, CSAIL

SUBMISSION DETAILS

Presentation Type Either Poster or Oral Presentation

Presentation Abstract Summary In blind individuals, traditional visual cortical areas are responsive to a range of nonvisual tasks, suggesting a functional reorganization in response to sensory loss. The general principles governing the dynamics of reorganized cortical processing remain unclear. Here we presented blind participants with braille characters while recording brain responses with MEG, then decoded the signals using multivariate pattern classification analysis. We found that letter identity was most readily decodable in sensorimotor cortex contralateral to the braille-stimulated finger, peaking ~125-200 ms after stimulus onset. Early visual and fusiform regions displayed more sustained decoding responses. The similarity structure of early MEG responses suggests an organizational scheme responsive to low-level features of the stimuli, i.e., the number of dots each character comprises, a scheme that shifted as the trial evolved. Our results could not only inform computational models of tactile processing but also expand future models of cortical processing optimized solely for visual stimuli to include crossmodal responses.

Paper Upload (PDF) [Teng_CCN_2017.pdf](#)

Co-author Information

* Presenting Author

First Name	Last Name	Affiliation	E-mail
Santani *	Teng *	Massachusetts Institute of Technology, CSAIL	santani@mit.edu
Radoslaw	Cichy	Freie University Berlin	rmcichy@zedat.fu-berlin.de
Dimitrios	Pantazis	MIT	pantazis@mit.edu
Aude	Oliva	MIT	oliva@mit.edu

Keywords

Keywords
blindness
braille
representational similarity analysis
multivariate pattern analysis (MVPA)