

# A Data-Driven Method for Investigating Cortical Functional Organization

**Submission ID** 3000331  
**Submission Type** Poster  
**Topic** Neuroscience  
**Status** Submitted  
**Submitter** Jason Webster  
**Affiliation** University of Washington

## SUBMISSION DETAILS

**Presentation Type** Either Poster or Oral Presentation

**Presentation Abstract Summary** We describe a novel cortical parcellation method that identifies regions with coherent functional response profiles, ‘grouping by response similarity’ (GRS). The algorithm does not require assumptions about stimulus properties, spatial relationships, or response uniformity. From fMRI responses to naturalistic videos in the human ventral temporal cortex, GRS finds discrete patches on the cortical surface with distinct functional response profiles. These regions show consistent boundaries across different stimulus sets, demonstrating that parcellation is not stimulus specific. Subsets of these regions correspond to the previously identified category-selective areas. Thus, grouping by response similarity provides a powerful exploratory analysis method for studying cortical organization in regions of cortex where functional roles are not yet clearly known.

**Paper Upload (PDF)** [CCN\\_paper\\_WebsterFine\\_20170529.pdf](#)

## Co-author Information

\* Presenting Author

First Name	Last Name	Affiliation	E-mail
Jason *	Webster *	University of Washington	jwebst@uw.edu
Ione	Fine	University of Washington	ionefine@uw.edu

## Keywords

Keywords
(f)MRI

parcellation

functional organization